



University of Groningen

Synthese en chemie van 4-azahomoadamantan-5-on

Korsloot, Jacob Gerard

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

1969

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Korsloot, J. G. (1969). Synthese en chemie van 4-azahomoadamantan-5-on. Groningen: s.n.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.


Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

SUMMARY

This thesis deals with the synthesis of 4-azatricyclo-[4.3.1.1^{3,8}]undecan-5-one (4-azahomoadamantan-5-one) and derivatives thereof. Moreover a new and convenient synthesis of 4-hydroxyadamantan-2-one is described.

After an introductory chapter, a review of the literature on variations of the adamantane skeleton and on the pharmacological and anti-viral properties of adamantane derivatives is given in chapter two.


The synthesis of 4-azahomoadamantan-5-one is the subject of chapter three. The Beckmann rearrangement of adamantanone oxime was shown to be a very complicated reaction, the product composition depending on the catalyst employed. With polyphosphoric acid the expected 4-azatricyclo[4.3.1.1^{3,8}]undecan-5-one was mainly obtained. With hydrogen chloride in acetonitrile adamantanone oxime also underwent rearrangement to the lactam, but in addition a fragmentation known as the "Beckmann fission" was observed. In this way bicyclo[3.3.1]non-6-ene-3-carbonitrile was obtained. Surprisingly a Ritter reaction of this unsaturated nitrile in concentrated sulfuric acid yielded 4-hydroxyadamantan-2-one as the main product. Isomerisation of the secondary hydroxy compound in the strongly acidic medium did not occur. A still more convenient synthesis of 4-hydroxyadamantan-2-one was possible by treating adamantanone oxime itself with concentrated sulfuric acid. The main reactions are shown in scheme I.



In chapter four the synthesis of derivatives of 4-azahomoadamantan-5-one is described. Four types of products were prepared as is shown in scheme II:

1. alkylated or acylated 4-azahomoadamantan-5-ones.
2. substituted 5-imino-4-azahomoadamantanes.
3. alkylated 4-azahomoadamantanes.
4. heterocycles fused with 4-azahomoadamantane.

SCHEME II



Furthermore the first synthesis of a diazadihomoadamantane was achieved by a Beckmann rearrangement of 4-azahomoadamantan-5-one oxime.

Finally the results of an investigation about the pharmacological and anti-viral properties of the synthesized compounds are given in chapter five. It was found that some of the compounds showed anti-viral and/or hypotensive activities.